NOVEL TYPE OF PHASE TRANSITION IN A SYSTEM OF SELF-DRIVEN PARTICLES

Tamás Vicsek, a,b András Czirók,
 a Eshel Ben-Jacob, c Inon Cohen,
 c

and Ofer Shochet, c

 ^a Department of Atomic Physics, Eötvös University, Budapest, Puskin u 5-7, 1088 Hungary

^b Institute for Technical Physics, Budapest, P.O.B. 76, 1325 Hungary

^c School of Physics, Tel-Aviv University, 69978 Tel-Aviv, Israel

Abstract

A simple model with a novel type of dynamics is introduced in order to investigate the emergence of self-ordered motion in systems of particles with biologically motivated interaction. In our model particles are driven with a constant absolute velocity and at each time step assume the average direction of motion of the particles in their neighborhood with some random perturbation (η) added. We present numerical evidence that this model results in a kinetic phase transition from no transport (zero average velocity, $|\mathbf{v}_a| = 0$) to finite net transport through spontaneous symmetry breaking of the rotational symmetry. The transition is continuous since $|\mathbf{v}_a|$ is found to scale as $(\eta_c - \eta)^\beta$ with $\beta \simeq 0.45$.

One of the most interesting aspects of many particle systems is that they exhibit a complex cooperative behavior during phase transition [1]. This remarkable feature of equilibrium systems has been studied in great detail for the last couple of decades leading to a deeper understanding of processes which may take place in an assembly of interacting particles. Concepts like scaling, universality and renormalization have resulted in a systematic picture of a wide range of systems in physics [1,2].

Recently there has been an increasing interest in the rich behaviour of systems which are far from equilibrium. Processes such as aggregation, viscous flows or biological pattern formation have been shown to involve scaling of the related geometrical and dynamic quantities characterizing these phenomena [3,4]. As a further similarity with equilibrium systems, the existence of phase transition type behavior has also been demonstrated in several investigations of growth processes [5-8]. These analogies with the basic features of equilibrium systems have represented a particularly important contribution to the understanding of the complex behavior of non-equilibrium processes.

In this work we introduce a model with a novel type of dynamics in order to investigate clustering, transport and phase transition in non-equilibrium systems where the velocity of the particles is determined by a simple rule and random fluctuations. The only rule of the model is: at each time step a given particle driven with a constant absolute velocity assumes the average direction of motion of the particles in its neighborhood of radius r with some random perturbation added. We show using simulations that in spite of its simplicity, this model results in a rich, realistic dynamics, including a kinetic phase transition from no transport to finite net transport through spontaneous symmetry breaking of the rotational symmetry.

In this sense our model is a *transport related*, *non-equilibrium analog of the ferromagnetic type of models*, with the important difference that it is inherently dynamic: the *elementary event is the motion of a particle* between two time steps. Thus, the analogy can be formulated as follows: the rule corresponding to the ferromagnetic interaction tending to align the spins in the same direction in the case of equilibrium models is replaced by the rule of *aligning the direction of motion* of particles in our model of cooperative motion. The level of random perturbations we apply are in analogy with the temperature.

Beyond the above aspects (analogies and simplicity) the proposed model is interesting because of possible applications in a wide range of biological systems involving clustering and migration. Biological subjects have the tendency to move as other subjects do in their neighborhood [9]. In addition to such trivial examples as schools of fish, herds of quadrupeds or flocks of flying birds, our model can be applied to the less known phenomena during bacterial colony growth [10]. There are bacteria (e.g., a strain of *Bacillus Subtilis*) which exhibit cooperative motion in order to survive under unfavorable conditions. The present model with some modifications is already capable to reproduce the main observed features of the motion (collective roatation and flocking) of bacteria [10]. Other biologically motivated, recent theoretical investigations of clustering, aggregation and orientational order in systems with diffusing directed objects have concentrated on the possible spatial patterns arising from an integro-differential equation approach and from cellular automata type models [11].

Furthermore, we expect that our model can be used to interpret the results of experiments on clustering and convection in a system of disks floating on an air table [12]. These experiments represent a physically motivated possible application of the present model, since they are being carried out in order to understand the flow of granular materials under specific conditions. We are aware that two groups are working on developing models similar to ours in order to interpret these air table experiments [13].

The actual simulations were carried out in a square shaped cell of linear size L with periodic boundary conditions. The particles were represented by points moving continuously (off-lattice) on the plane. We used the interaction radius r as unit to measure distances (r = 1) while the time unit, $\Delta t = 1$ was the time interval between

two updating of the directions/positions. In most of our simulations we used the simplest initial conditions: i) at time t = 0 N particles were randomly distributed in the cell and ii) had the same absolute velocity v and iii) randomly distributed directions θ . The velocities $\{\mathbf{v}_i\}$ of the particles were determined simultaneously at each time step and the position of the *i*th particle updated according to

$$\mathbf{x}_i(t+1) = \mathbf{x}_i(t) + \mathbf{v}_i(t)\Delta t.$$
(1)

Here the velocity of a particle $\mathbf{v}_i(t+1)$ was constructed to have an absolute value v and a direction given by the angle $\theta(t+1)$. This angle was obtained from the expression

$$\theta(t+1) = \langle \theta(t) \rangle_r + \Delta \theta, \tag{2}$$

where $\langle \theta(t) \rangle_r$ denotes the average direction of the velocities of particles (including particle *i*) being within a circle of radius *r* surrounding the given particle. The average direction was given by the angle $\operatorname{arctg}[\langle \sin(\theta(t)) \rangle_r / \langle \cos(\theta(t)) \rangle_r]$. In (2) $\Delta \theta$ is a random number chosen with a unifom probability from the interval $[-\eta/2, \eta/2]$. Thus, the term $\Delta \theta$ represents noise which we shall use as a temperature-like variable. Correspondingly, there are three free parameters for a given system size: η , ρ and v, where v is the distance a particle makes between two updatings.

We have chosen this realization because of its simplicity, however, there may be several more interesting alternatives of implementing the main rules of the model. In particular, the absolute value of the velocities does not have to be fixed, one can introduce further kind of interaction between a particles and or consider lattice alternatives of the model. In the rest of this paper we shall concentrate on the above described simplest version and investigate the nontrivial behavior of the transport properties as the two basic parameters of the model, the noise η and the density $\rho = N/L^2$ are varied. We used v = 0.03 in the simulations we report on because of the following reasons. In the limit $v \to 0$ the particles do not move and the model becomes an analog of the well known XY model. For $v \to \infty$ the particles become completely mixed between two updates and this limit corresponds to the so called mean field behavior of a ferromagnet. We use v = 0.03 for which the particles always interact with their actual neighbours and move fast enough to change the configuration after a few updates of the directions. According to our simulations in a wide range of the velocities (0.003 < v < 0.3) the actual value of v does not affect the results.

Fig. 1a-d demonstrates the velocity fields during runs with various selections for the value of the parameters ρ and η . The actual velocity of a particle is indicated by a small arrow, while their trajectory for the last 20 time steps is shown by a short continuous curve. (a) At t = 0 the positions and the direction of velocities are distributed randomly. (b) For small densities and noise the particles tend to form groups moving coherently in random directions. (c) At higher densities and noise the particles move randomly with some correlation. (d) perhaps the most interesting case is when the density is large and the noise is small; in this case the motion becomes ordered on a macroscopic scale and all of the particles tend to *move in the same spontaneously selected direction*.

This kinetic phase transition is due to the fact that the particles are driven with a constant absolute velocity; thus, unlike standard physical systems in our case the *net momentum of the interacting particles is not conserved* during collision. We have studied in detail the nature of this transition by determining the absolute value of the average normalized velocity

$$v_a = \frac{1}{Nv} \left| \sum_{i=1}^{N} \mathbf{v}_i \right| \tag{3}$$

of the entire system of particles as the noise and the density were changed. This velocity is approximately zero if the direction of the motion of the individual particles is distributed randomly, while for the coherently moving phase (with ordered direction of velocities) $v_a \simeq 1$, so that we can consider the average velocity as an *order parameter*.

First we gradually decreased the amount of noise η in cells of various sizes for a fixed density ρ and observed a transition from a disorderly moving phase to a phase with a coherent motion of the particles (Fig. 2a). The uncertainity of the data points is within the range of the symbols except for runs carried out with 4000 and 10000 particles close to the transition. For these η values the statistical errors estimated from 5 runs with different initial conditions are in the range of 5% (resulting in an overlap of the results for a limited number of η values) due to the slow convergence and large fluctuations. In Fig. 2b we show how v_a changes if the noise is kept constant and the density is increased.

Quite remarkably, the behavior of the kinetic order parameter v_a is very similar to that of the order parameter of some equilibrium systems close to their critical point. The strongest indication of a transition in our nonequilibrium model is the fact that as we go to larger system sizes the region over which the data show scaling is increasing (see Fig. 3a). Only an extremely unusual crossover could change this *tendency*. A plausible physical picture behind our finding is the following: since the particles are diffusing, there is mixing in the system resulting in an effective (long range) interaction radius.

Thus, we can assume that in the thermodynamic limit our model exhibits a kinetic phase transition analogous to the continuous phase transition in equilibrium systems, i.e.,

$$v_a \sim (\eta_c(\rho) - \eta)^{\beta}$$
 and $v_a \sim (\rho - \rho_c(\eta)),^{\delta}$ (4)

where β and δ are critical exponents and $\eta_c(\rho)$ and $\rho_c(\eta)$ are the critical noise and density (for $L \to \infty$), respectively. We can determine β and δ corresponding to the rate of vanishing of the order parameter from plotting $\log v_a$ as a function of $\log[\eta_c(L) - \eta)/\eta_c(L)]$ and $\log[\rho - \rho_c(L)/\rho_c(L)]$ for some fixed values of ρ and η , respectively (Fig. 3). For finite sizes $\eta_c(L)$ and $\rho_c(L)$ are L dependent, thus, we used such values of quantities for which the plots in Fig. 3 were the straightest in the relevant region of noise or density values. The slope of the lines fitted to the data can be associated with the critical exponents for which we obtained $\beta = 0.45 \pm 0.07$ and $\delta = 0.35 \pm 0.06$. The errors in determining β and δ are due to the uncertainties in the i) v_a and the ii) $\eta_c(L)$ and $\rho_c(L)$ values. Since the scaling plots in Fig. 3 depend sensitively on the choice of the critical noise and density and our method of determing their value is indirect (from the straightness of the data sets) we give rather conservative estimates for the errors of β and δ .

We have carried out a finite size scaling analysis of $\eta_c(L)$ and obtained $\eta_c(\infty) = 2.9 \pm 0.05$ for $\rho = 0.4$ (note that the "infinite temperature" limit of our model is $\eta_c = 2\pi$). As indicated, η_c depends on ρ , in fact, we expect a phase diagram (a line of critical temperatures) analogous to that of disordered ferromagnets, η_c playing the role of temperature and ρ playing the role of the density of spins. In this case β and δ are expected to have the same value. On the other hand, strong crossover effects are likely to effect their actual values in a finite size simulation. Although our estimates for β and δ are different, on the basis of our simulations we cannot exclude the possibility (allowed by our error bars) that they become equal in the thermodynamic limit. However, the determination of the phase diagram and a more precise calculation of the exponents of our new model is outside of the scope of the present work which concentrates on demonstrating the main features of a novel nonequilibrium system.

The emergence of cooperative motion in our model has analogies with the appearance of spatial order in equilibrium systems. This fact, and the simplicity of our model suggests that, with appropriate modifications, the theoretical methods for describing critical phenomena may be applicable to the present kind of far-fromequilibrium phase transition. The kinetic phase transitions which have been observed in surface growth models [5-8] are both in analogy and different from the situation described here. The similarity is in the scaling behaviour of an inherently non-equilibrium order parameter, while the two kinds of processes are distinct from the point of the driving force acting on the particles. Self-driven particles are uncommon in physics, but they are typical in biological systems, including live organisms and the so called "molecular motors" having attracted great interest recently [14]. Transitions have been observed in traffic models [15] consisting of particles (cars) which can also be interpreted as self-driven particles.

There are interesting further variations of the model investigated in this work. It is expected that taking into account a hard core term in the interaction or using semi periodic or open boundary conditions results in additional non-trivial effects. Our preliminary results [10] indicate that a model with hard core repulsion and specific boundary conditions can be successfully used to interpret recent observations of coherent motions in geometrically complex bacterial colonies growing on soft agar surfaces [10,16-18].

Acknowledgement: The present research in part was supported by the Hungarian Research Foundation Grant No. T4439, by a grant from the German-Israeli Foundation for Scientific Research and Development, and by the Program for Alternative Thinking at Tel-Aviv University.

References

- H. E. Stanley Introduction to Phase Transitions and Critical Phenomena (Oxford University Press, Oxford, 1971)
- [2] see, e.g., S.-K. Ma Statistical Mechanics (World Scientific, Singapore, 1985); Modern Theory of Critical Phenomena (Benjamin, New York, 1976)
- [3] P. Meakin, in *Phase Transitions and Critical Phenomena* Vol 12. edited by C.
 Domb and J. Lebowitz (Academic Press, New York, 1987)
- [4] T. Vicsek, Fractal Growth Phenomena. (Singapore, World Scientific, 1992)
- [5] R. Jullien, J. Kertész, P. Meakin and D. Wolf, eds., Surface Disordering. (New York, Nova Science, 1992)
- [6] J. Kertész, and D. E. Wolf, Phys. Rev. Lett. 62, (1989) 2571
- [7] N. Martys, M. Cieplak, and M. O. Robbins, *Phys. Rev. Lett.* 66, (1991) 1058
- [8] Z. Csahok, K. Honda, E. Somfai, M. Vicsek and T. Vicsek, *Physica* A200, (1993)
 136
- [9] D. P. O'Brien J. Exp. Mar. Biol. Ecol. **128**, (1989) 1
- [10] E. Ben-Jacob, A. Czirók, I. Cohen, O. Shochet, A. Tenenbaum and T. Vicsek to be published
- [11] L. Edelstein-Keshet and G. B. Ermentrout, J. Math. Biol. 29, (1990) 33
- [12] J. Lemaitre, A. Gervois, H. Peerhossaini, D. Bideau and J. P. Troadec, J. Phys.
 D: Appl. Phys. 23 (1990) 1396
- [13] H. J. Herrmann, private communication
- [13] see, e.g., R. D. Astumian and M. Bier Phys. Rev. Lett. 72, (1994) 1766
- [14] see, e.g., K. Nagel and H. J. Herrmann, Physica A 199, (1993) 254 and references therein
- [15] E. Ben-Jacob, H. Shmueli, O. Shochet and A. Tenenbaum *Physica* A187, (1992) 378
- [16] E. Ben-Jacob, I. Cohen, O. Shochet, A. Tenenbaum, A. Czirók, and T. Vicsek,

Nature 368, 46 (1994)

[17] T. Matsuyama, R. M. Harshey and M. Matsushita, $Fractals~{\bf 1},~302~(1993)$

Figure captions

Figure 1. In this figure the velocities of the particles are displayed for varying values of the density and the noise. The actual velocity of a particle is indicated by a small arrow, while their trajectory for the last 20 time step is shown by a short continuous curve. The number of particles is N = 300 in each case. (a) t = 0, L = 7 $\eta = 2.0$ (b) For small densities and noise the particles tend to form groups moving coherently in random directions, here L = 25, $\eta = 0.1$; (c) After some time at higher densities and noise (L = 7 $\eta = 2.0$) the particles move randomly with some correlation; (d) For higher density and *small noise* (L = 5 $\eta = 0.1$) the motion becomes ordered. All of our results shown in Figs. 1-3 were obtained from simulations in which v was set to be equal to 0.03.

Figure 2. (a) The absolute value of the average velocity (v_a) versus the noise η in cells of various sizes for a fixed density ρ . The symbols correspond to $\Box - N = 40, L = 3.1;$ $+ - N = 100, L = 5; \times - N = 400, L = 10; \triangle - N = 4000, L = 31.6. \diamondsuit - N = 10000,$ L = 50. In Fig. 2b (for L = 20) we show how v_a changes if the noise is kept constant and the density is increased.

Fig. 3 Dependence of $\log v_a$ on $\log[(\eta_c(L) - \eta)/\eta_c(L)]$ and $\log[(\rho - \rho_c(L)/\rho_c(L)]]$. The slope of the lines fitted to the data can be associated with the critical exponents β and δ . a) is for $\rho = 0.4$, b) is for L = 20 and $\eta = 2.0$











$$\eta$$



