

3.3 Models, Validity, and Satisfiability

F is *valid* in \mathcal{A} under assignment β :

$$\mathcal{A}, \beta \models F \quad :\Leftrightarrow \quad \mathcal{A}(\beta)(F) = 1$$

F is *valid* in \mathcal{A} (\mathcal{A} is a *model* of F):

$$\mathcal{A} \models F \quad :\Leftrightarrow \quad \mathcal{A}, \beta \models F, \text{ for all } \beta \in X \rightarrow U_{\mathcal{A}}$$

F is *valid* (or is a *tautology*):

$$\models F \quad :\Leftrightarrow \quad \mathcal{A} \models F, \text{ for all } \mathcal{A} \in \Sigma\text{-Alg}$$

F is called *satisfiable* iff there exist \mathcal{A} and β such that $\mathcal{A}, \beta \models F$. Otherwise F is called *unsatisfiable*.

Substitution Lemma

The following propositions, to be proved by structural induction, hold for all Σ -algebras \mathcal{A} , assignments β , and substitutions σ .

Lemma 3.3 For any Σ -term t

$$\mathcal{A}(\beta)(t\sigma) = \mathcal{A}(\beta \circ \sigma)(t),$$

where $\beta \circ \sigma : X \rightarrow \mathcal{A}$ is the assignment $\beta \circ \sigma(x) = \mathcal{A}(\beta)(x\sigma)$.

Proposition 3.4 For any Σ -formula F , $\mathcal{A}(\beta)(F\sigma) = \mathcal{A}(\beta \circ \sigma)(F)$.

Corollary 3.5 $\mathcal{A}, \beta \models F\sigma \Leftrightarrow \mathcal{A}, \beta \circ \sigma \models F$

These theorems basically express that the syntactic concept of substitution corresponds to the semantic concept of an assignment.

Entailment and Equivalence

F entails (implies) G (or G is a consequence of F), written $F \models G$, if for all $\mathcal{A} \in \Sigma\text{-Alg}$ and $\beta \in X \rightarrow U_{\mathcal{A}}$, whenever $\mathcal{A}, \beta \models F$, then $\mathcal{A}, \beta \models G$.

F and G are called *equivalent*, written $F \models\!\!\!\!\!\! \models G$, if for all $\mathcal{A} \in \Sigma\text{-Alg}$ und $\beta \in X \rightarrow U_{\mathcal{A}}$ we have $\mathcal{A}, \beta \models F \Leftrightarrow \mathcal{A}, \beta \models G$.

Proposition 3.6 F entails G iff $(F \rightarrow G)$ is valid

Proposition 3.7 F and G are equivalent iff $(F \leftrightarrow G)$ is valid.

Extension to sets of formulas N in the “natural way”, e. g., $N \models F$

$:\Leftrightarrow$ for all $\mathcal{A} \in \Sigma\text{-Alg}$ and $\beta \in X \rightarrow U_{\mathcal{A}}$: if $\mathcal{A}, \beta \models G$, for all $G \in N$, then $\mathcal{A}, \beta \models F$.

Validity vs. Unsatisfiability

Validity and unsatisfiability are just two sides of the same medal as explained by the following proposition.

Proposition 3.8 Let F and G be formulas, let N be a set of formulas. Then

- (i) F is valid if and only if $\neg F$ is unsatisfiable.
- (ii) $F \models G$ if and only if $F \wedge \neg G$ is unsatisfiable.
- (iii) $N \models G$ if and only if $N \cup \{\neg G\}$ is unsatisfiable.

Hence in order to design a theorem prover (validity checker) it is sufficient to design a checker for unsatisfiability.

Theory of a Structure

Let $\mathcal{A} \in \Sigma\text{-Alg}$. The (*first-order*) *theory* of \mathcal{A} is defined as

$$Th(\mathcal{A}) = \{ G \in F_{\Sigma}(X) \mid \mathcal{A} \models G \}$$

Problem of axiomatizability:

For which structures \mathcal{A} can one *axiomatize* $Th(\mathcal{A})$, that is, can one write down a formula F (or a recursively enumerable set F of formulas) such that

$$Th(\mathcal{A}) = \{ G \mid F \models G \}?$$

Analogously for sets of structures.

Two Interesting Theories

Let $\Sigma_{Pres} = (\{0/0, s/1, +/2\}, \emptyset)$ and $\mathbb{Z}_+ = (\mathbb{Z}, 0, s, +)$ its standard interpretation on the integers. $Th(\mathbb{Z}_+)$ is called *Presburger arithmetic* (M. Presburger, 1929). (There is no essential difference when one, instead of \mathbb{Z} , considers the natural numbers \mathbb{N} as standard interpretation.)

Presburger arithmetic is decidable in 3EXPTIME (D. Oppen, JCSS, 16(3):323–332, 1978), and in 2EXPSPACE, using automata-theoretic methods (and there is a constant $c \geq 0$ such that $Th(\mathbb{Z}_+) \notin \text{NTIME}(2^{2^{cn}})$).

However, $\mathbb{N}_* = (\mathbb{N}, 0, s, +, *)$, the standard interpretation of $\Sigma_{PA} = (\{0/0, s/1, +/2, */2\}, \emptyset)$, has as theory the so-called *Peano arithmetic* which is undecidable, not even recursively enumerable.

Note: The choice of signature can make a big difference with regard to the computational complexity of theories.

3.4 Algorithmic Problems

Validity(F): $\models F$?

Satisfiability(F): F satisfiable?

Entailment(F, G): does F entail G ?

Model(A, F): $A \models F$?

Solve(A, F): find an assignment β such that $A, \beta \models F$.

Solve(F): find a substitution σ such that $\models F\sigma$.

Abduce(F): find G with “certain properties” such that $G \models F$.

Gödel’s Famous Theorems

1. For most signatures Σ , validity is undecidable for Σ -formulas. (One can easily encode Turing machines in most signatures.)
2. For each signature Σ , the set of valid Σ -formulas is recursively enumerable. (We will prove this by giving complete deduction systems.)
3. For $\Sigma = \Sigma_{PA}$ and $\mathbb{N}_* = (\mathbb{N}, 0, s, +, *)$, the theory $Th(\mathbb{N}_*)$ is not recursively enumerable.

These complexity results motivate the study of subclasses of formulas (*fragments*) of first-order logic

Q: Can you think of any fragments of first-order logic for which validity is decidable?

Some Decidable Fragments

Some decidable fragments:

- *Monadic class*: no function symbols, all predicates unary; validity is NEXPTIME-complete.
- Variable-free formulas without equality: satisfiability is NP-complete. (why?)
- Variable-free Horn clauses (clauses with at most one positive atom): entailment is decidable in linear time.
- Finite model checking is decidable in time polynomial in the size of the structure and the formula.

3.5 Normal Forms and Skolemization

Study of normal forms motivated by

- reduction of logical concepts,
- efficient data structures for theorem proving.

The main problem in first-order logic is the treatment of quantifiers. The subsequent normal form transformations are intended to eliminate many of them.

Prenex Normal Form

Prenex formulas have the form

$$Q_1x_1 \dots Q_nx_n F,$$

where F is quantifier-free and $Q_i \in \{\forall, \exists\}$; we call $Q_1x_1 \dots Q_nx_n$ the *quantifier prefix* and F the *matrix* of the formula.

Computing prenex normal form by the rewrite relation \Rightarrow_P :

$$\begin{aligned} (F \leftrightarrow G) &\Rightarrow_P (F \rightarrow G) \wedge (G \rightarrow F) \\ \neg Qx F &\Rightarrow_P \overline{Q}x \neg F && (\neg Q) \\ ((Qx F) \rho G) &\Rightarrow_P Qy(F\{x \mapsto y\} \rho G), \rho \in \{\wedge, \vee\} \\ ((Qx F) \rightarrow G) &\Rightarrow_P \overline{Q}y(F\{x \mapsto y\} \rightarrow G), \\ (F \rho (Qx G)) &\Rightarrow_P Qy(F \rho G\{x \mapsto y\}), \rho \in \{\wedge, \vee, \rightarrow\} \end{aligned}$$

Here y is always assumed to be some fresh variable and \overline{Q} denotes the quantifier *dual* to Q , i. e., $\overline{\forall} = \exists$ and $\overline{\exists} = \forall$.

Skolemization

Intuition: replacement of $\exists y$ by a concrete choice function computing y from all the arguments y depends on.

Transformation \Rightarrow_S (to be applied outermost, *not* in subformulas):

$$\forall x_1, \dots, x_n \exists y F \Rightarrow_S \forall x_1, \dots, x_n F \{y \mapsto f(x_1, \dots, x_n)\}$$

where f/n is a new function symbol (*Skolem function*).

Together: $F \xRightarrow{*}_P \underbrace{G}_{\text{prenex}} \xRightarrow{*}_S \underbrace{H}_{\text{prenex, no } \exists}$

Theorem 3.9 Let F , G , and H as defined above and closed. Then

- (i) F and G are equivalent.
- (ii) $H \models G$ but the converse is not true in general.
- (iii) G satisfiable (w. r. t. Σ -Alg) $\Leftrightarrow H$ satisfiable (w. r. t. Σ' -Alg) where $\Sigma' = (\Omega \cup SKF, \Pi)$, if $\Sigma = (\Omega, \Pi)$.

Clausal Normal Form (Conjunctive Normal Form)

$$\begin{aligned} (F \leftrightarrow G) &\Rightarrow_K (F \rightarrow G) \wedge (G \rightarrow F) \\ (F \rightarrow G) &\Rightarrow_K (\neg F \vee G) \\ \neg(F \vee G) &\Rightarrow_K (\neg F \wedge \neg G) \\ \neg(F \wedge G) &\Rightarrow_K (\neg F \vee \neg G) \\ \neg\neg F &\Rightarrow_K F \\ (F \wedge G) \vee H &\Rightarrow_K (F \vee H) \wedge (G \vee H) \\ (F \wedge \top) &\Rightarrow_K F \\ (F \wedge \perp) &\Rightarrow_K \perp \\ (F \vee \top) &\Rightarrow_K \top \\ (F \vee \perp) &\Rightarrow_K F \end{aligned}$$

These rules are to be applied modulo associativity and commutativity of \wedge and \vee . The first five rules, plus the rule $(\neg Q)$, compute the *negation normal form* (NNF) of a formula.

The Complete Picture

$$\begin{aligned}
 F &\xRightarrow{*}_P Q_1 y_1 \dots Q_n y_n G && (G \text{ quantifier-free}) \\
 &\xRightarrow{*}_S \forall x_1, \dots, x_m H && (m \leq n, H \text{ quantifier-free}) \\
 &\xRightarrow{*}_K \underbrace{\underbrace{\forall x_1, \dots, x_m}_{\text{leave out}} \bigwedge_{i=1}^k \underbrace{\bigvee_{j=1}^{n_i} L_{ij}}_{\text{clauses } C_i}}_{F'}
 \end{aligned}$$

$N = \{C_1, \dots, C_k\}$ is called the *clausal (normal) form* (CNF) of F .

Note: the variables in the clauses are implicitly universally quantified.

Theorem 3.10 *Let F be closed. Then $F' \models F$. (The converse is not true in general.)*

Theorem 3.11 *Let F be closed. Then F is satisfiable iff F' is satisfiable iff N is satisfiable*

Optimization

The normal form algorithm described so far leaves lots of room for optimization. Note that we only can preserve satisfiability anyway due to Skolemization.

- size of the CNF is exponential when done naively; the transformations we introduced already for propositional logic avoid this exponential growth;
- we want to preserve the original formula structure;
- we want small arity of Skolem functions (see next section).